

ORIGINAL STUDY

Tomographic dimensions of the nasal vestibular body in relation to nasal obstruction in adults

Luis Gerardo Vargas-Cárdenas¹, Fernando Juan Ramírez-Oropeza², Omar Gómez-Monterrosas³, Juan Antonio Lugo-Machado⁴ 

¹Department of Otolaryngology and Head and Neck Surgery and Facial Surgery, Otorrino Autlan Gerardo Vargas Cardenas, Autlan, Jalisco, Mexico

²Otolaryngology and Facial Surgery Department, National Autonomous University of Mexico, Puebla, Mexico

³Cardiology Service. Ángeles Hospital, Puebla, Mexico

⁴Otolaryngology Department, "Lic. Luis Donaldo Colosio Murrieta" Specialty Hospital Number 2, IMSS, Ciudad Obregon, Sonora, Mexico

ABSTRACT

INTRODUCTION. The nasal vestibular body is a recently reported dynamic elevation in the vicinity of the inferolateral internal nasal valve. The contribution of this anatomical zone in persistent nasal obstruction is not yet defined.

OBJECTIVE. To assess the correlation between the tomographic size of the nasal vestibular body and the presence of nasal obstruction in adults.

MATERIAL AND METHODS. An observational, analytical, prospective, cross-sectional study of a series of cases was conducted between August 1 and December 30, 2020. We included 23 adult patients and we recorded the sex, age, symptoms of obstruction, presence of allergic rhinitis, whether they used treatment for rhinitis, presence of septal deviation and dimensions in length and width of the nasal vestibular body. Patients were distributed in groups, with and without nasal obstruction, and the area of the vestibular body was measured in digital format using millimetre computed tomography.

RESULTS. Of a total of 23 cases, 10 were men (43.48%) and 13 women (56.52%), mean age 28.8±10.2 years. 16 patients (69.57%) had symptoms of nasal obstruction and 7 (30.43%) were without obstruction. 10 patients (43.48%) presented symptoms of rhinitis. Using Spearman's correlation test, we found a positive correlation between width ($R = 0.79$, $p < 0.001$) and nasal vestibular body length ($R = 0.77$, $p < 0.001$) and nasal obstruction.

CONCLUSION. The nasal vestibular body is a structure that could be related to the symptoms of nasal obstruction in the adult population.

KEYWORDS: nasal obstruction, tomography, nasal vestibular body.

INTRODUCTION

Among the critical components of nasal respiratory physiology, the nasal valve area plays an important role in terms of airflow through the nasal cavity during inspiration, but also in the subjective perception of nasal breathing by each individual¹. Having et al. has already studied the different physical aspects that play an important role in this area, such as the Bernoulli's princi-

ple, the Venturi effect or the Poiseuille's law, the harmonic and dynamic function of the nasal valves and their different structures that compose them, which establishes a dynamic at the entrance of the upper airways².

The nasal vestibular body (NVB) has been described since 1971 in some mammalian species³. However, its implications in persistent nasal obstruction in humans have only recently been reported, given its proximity to the nasal valve⁴.

Corresponding author: Juan Antonio Lugo Machado, Address: Calle Hidalgo Pte. S/N, Bellavista, 85130 Ciudad Obregón, Mexico

ORCID: <https://orcid.org/0000-0003-4864-8546>

e-mail: juan.lugo.imss@gmail.com

Received for publication: 18 October 2021 / **Accepted:** 29 November 2021




Figure 1. Illustrating the nasal vestibular body (in pink) and the nasal septal body (in black).

Clinically, the evaluation of symptoms of nasal obstruction focuses largely on the direct visualization of the nasal cavity through the anterior rhinoscopy^{5,6}. Different tests have been used to objectively quantify nasal obstruction, such as acoustic rhinometry⁶, identification of the maximum nasal inflammatory flow⁷, or validated subjective questionnaires^{8,9}. However, there is no consensus in their use in establishing the correct diagnosis.

Anatomically, NVB arises from the infero-lateral edge of the nose, between the vestibule and the region of the internal nasal valve, and it is adjacent to the anterior part of the inferior turbinate (Figure 1). Therefore, it is reasonable that the nasal vestibular body may be a component in the nasal obstruction experienced by some patients¹. The low notoriety of this structure makes it very common for it to be easily overlooked¹.

If the implication of this anatomical area in a decrease in nasal airflow is not recognized, it may compromise the results of standard surgical interventions for nasal obstruction. It is very important not to confuse the NVB with the nasal septal body which is a different structure, located in the anterior part of the nasal septum, adjacent to the anterior part of the middle turbinate and to the upper part of the lower turbinate^{10,11} (Figure 1).

At present, nasal obstruction is a topic that should be fully understood by the otolaryngologist, since mismanagement or exclusion of an anatomical site to be treated in surgery can be the cause of an unsatisfactory result, especially in the areas of the nasal valve, which can have a high impact on the final breathing. There is little international and national literature that addresses the diagnosis and imaging of this anatomical area of the nose¹¹.

This study focuses on measuring the nasal vestibular body, which is a recently described dynamic structure that has been the focus of different studies^{1,4}. It was observed that NVB could have an additional benefit in nasal ventilation, if this structure is surgically reduced. However, due to its recent description, there is no tool with which it can be measured or observed objectively, a situation that could serve in future surgical planning.

MATERIAL AND METHODS

An analytical, prospective, cross-sectional observational study of a consecutive series of cases, chosen in a non-probabilistic manner, was car-




Figure 2. Crano-facial CT scan, coronal slices, where the nasal vestibular body and the septal vestibular body are indicated.

ried out at the Facial Surgery and Cosmetology Clinic of Puebla, between August 1, 2020, and December 30, 2020.

The study included patients over 18 years of age, who came for a nasal aesthetic procedure, routinely asked to perform simple tomography of the nose and paranasal sinuses. Exclusion criteria: alterations in nasal anatomy (such as nasal fractures), nasal polyps, previous nasal surgeries, nasal tumors, cleft lip or palate.

Of a total of 32 patients evaluated, 23 were included in whom the variables age, sex, symptoms of nasal obstruction, the presence or absence of rhinitis, the use of topical intranasal medication, the size (in millimetres) of the NVB in length and width were registered. The patients included in the study were distributed into two groups: patients with nasal obstruction, patients without nasal obstruction. In all cases, a crano-facial CT scan was performed and the images were sent to an external collaborator specialized in radiology who analysed and measured the anatomical area with RadiAnt DICOM Viewer 2020 program (Figure 2).

RESULTS

In the study period from August 1 to December 30, 2020, 23 cases were included, 10 male patients (43.48%) and 13 women (56.52%), with a mean age of 28.8 ± 10.2 years (Table 1).

Symptoms of nasal obstruction were present in 16 patients (69.57%) while 7 (30.43%) had no nasal ob-

Table 1. Main demographic characteristics of the patients included in our study group.

Age (mean \pm SD)	Male	Female	Average
	33 +16.05	25.3 + 12.20	28.8 \pm 10.2
Gender	Male	Female	Total
	10 (43.48%)	13 (56.52%)	23

Table 2. Average of measurements in millimetres of the nasal vestibular body.

Average in mm	Height right	Length right	Height left	Length left
GENERAL	3.8	5.9	2.95	5.88
Obstruction	3.78	6.63	3.58	6.63
No obstruction	1.55	4.31	1.51	4.15
Rhinitis	4.2	6.96	4	6.8
No rhinitis	2.16	5.21	2.14	5.17
Treatment	3.96	6.76	3.75	6.56
No treatment	4.55	7.25	4.37	7.15

struction. 10 cases (43.48%) had symptoms of rhinitis. 13 patients (56.52%) did not use any medication and only 6 (26%) of them mentioned using some type of medication. Of the 10 patients who reported rhinitis problems, 9 reported nasal obstruction, while of the 12 patients who had some type of septal deviation on the CT scan, 10 reported nasal obstruction.

The average of the dimensions of the nasal vestibular body according to each clinical symptom are presented in Table 2. The table shows the general average of measurements of both the length and the height of both sides of the NVB. After analysing the data, we observed that the dimensions of the NVB are significantly higher in patients with nasal obstruction and rhinitis when compared with those who did not present this pathology.

Using the Spearman's correlation test, a moderate positive correlation was found between nasal obstruction and nasal vestibular body width ($R = 0.79$, $p < 0.001$), and nasal obstruction and nasal vestibular body length ($R = 0.77$, $p < 0.001$). However, there was only a weak and non-significant correlation between the degree of obstruction and patients who presented with rhinitis ($R = 0.39$, $p = 0.06$) (Graph 1).

Graph 1 shows the comparison of the 2 measurements made of both the length and the height of both sides' nasal vestibular bodies, observing a clear preponderance of higher volumes in patients who report nasal obstruction, as well as in patients who report rhinitis symptoms.

DISCUSSIONS


The nasal vestibular body is a structure recently considered as a factor of nasal obstruction. There are very limited publications on the implications related to this anatomical area. Authors such as Garrett D. Locketz et al.¹ refer, in 2016, to this structure as a cause of nasal obstruction. More recently, Nour Ibrahim et al.⁴ has published a clinical-surgical study concluding that the surgical treatment of the NVB can be highly effective in treating persistent or recalcitrant nasal obstruction.

Our results could correlate the dimensions of NVB with the symptoms of nasal obstruction. However, there are a number of limitations, the most important being the small number of patients included in the study, that does not allow the generalization of the results to a population sample. At the same time, publications at national and international level are very scarce.

We could point out that this short series represents the first publication that attempts to objectively measure and correlate the effect of the dimensions of this nasal structure and obstruction.

CONCLUSIONS

The dimensions of the nasal vestibular body can be measured by tomography, and this could be correlated with the symptoms of nasal obstruction.



Graph 1. Comparison of the nasal vestibular body measurements (mm), according to different symptoms.

We believe that it is necessary to carry out studies with a larger population, trying to control some variables that could generate biases and limitations in the results.

Contribution of author: (1) Concept or design: LGVC, (2) Data acquisition: LGVC, (3) Data analysis or interpretation: GMO, (4) Article writing: FJRO, (5) Critical review for important intellectual content: JALM.

Conflicts of interest: All authors have not disclosed any conflicts of interest

Acknowledgments: We thank the Hospital Alta Especialidad en Rinología y Cirugía Facial of the National Autonomous University of Mexico. This research has been presented as a thesis for a high specialty in rhinology and facial surgery endorsed by the National Autonomous University of Mexico.

Funding/Support: This research did not receive any specific grants from any funding agency in the public, commercial, or non-profit sectors.

Ethical approval: This project was submitted for evaluation by the ethics committee of the rhinology and facial surgery clinic in Puebla, Mexico.

REFERENCES

- Locketz GD, Teo NW, Walgama E, Humphreys IM, Nayak J V. The nasal vestibular body: anatomy, clinical features, and treatment considerations. *Eur Arch Otorhinolaryngol.* 2016;273(3):777-81.
- Daniel RK, Pálházi P. Rhinoplasty. *An Anatomical and Clinical Atlas.* Springer; 2018, p. 337.
- Bojsen-Moller F, Fahrenkrug J. Nasal swell-bodies and cyclic changes in the air passage of the rat and rabbit nose. *J Anat.* 1971;110(Pt 1):25-37.
- Ibrahim N, Tyler MA, Borchard NA, Rathor A, Nayak JV. Nasal vestibular body treatment for recalcitrant nasal obstruction. *Int Forum Allergy Rhinol.* 2020;10(3):388-94. DOI: 10.1002/alr.22463.
- Dastidar P, Numminem J, Hainonen T, Ryymin P, Rautiainen M, Laasonen E. Nasal airway volumetric measurement using segmented HRCT images and acoustic rhinometry. *Am J Rhinol.* 1999;13(2):97-103.
- Hilberg O, Jackson AC, Swift DL, Pedersen OF. Acoustic rhinometry: Evaluation of nasal cavity geometry by acoustic reflection. *J Appl Physiol* (1985). 1989;66(1):295-303. DOI: 10.1152/jappl.1989.66.1.295.
- Ganslmayer M, Spertini F, Rahm F, Terrien MH, Mosimann B, Leimgruber A. Evaluation of acoustic rhinometry in a nasal provocation test with allergen. *Allergy.* 1999;54(9):974-9.
- Lam DJ, James KT, Weaver EM. Comparison of anatomic, physiological, and subjective measures of the nasal airway. *Am J Rhinol.* 2006;20(5):463-70. DOI: 10.2500/ajr.2006.20.2940.
- Lipan MJ, Most SP. Development of a severity classification system for subjective nasal obstruction. *JAMA Facial Plast Surg.* 2013;15(5):358-61. DOI: 10.1001/jamafacial.2013.344.
- Wexler D, Braverman I, Amar M. Histology of the nasal septal swell body (septal turbinate). *Otolaryngol Head Neck Surg.* 2006;134(4):596-600. DOI: 10.1016/j.otohns.2005.10.058.
- Yu MS, Choi CH, Jung MS, Kim HC. Correlation between septal body size and inferior turbinate hypertrophy on computerised tomography scans in fifty patients: A radiological analysis. *Clin Otolaryngol.* 2018;43(3):952-5. DOI: 10.1111/coa.13073.

© 2022 Luis Gerardo Vargas-Cárdenas, Fernando Juan Ramírez-Oropeza, Omar Gómez -Monterrosas, Juan Antonio Lugo-Machado, published by Romanian Rhinologic Society



This is an open access article published under the terms and conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>). CC BY-NC-ND 4.0 license requires that reusers give credit to the creator by citing or quoting the original work. It allows reusers to copy, share, read, download, print, redistribute the material in any medium or format, or to link to the full texts of the articles, for non-commercial purposes only. If others remix, adapt, or build upon the material, they may not distribute the modified material.