

LITERATURE REVIEW**Rhinitis and sinusitis in the older population and its association with elderly asthma****Cemal Cingi¹**, **Nuray Bayar Muluk²**, **Arzu Yorgancıoğlu³**, **Necdet Demir⁴**, **Bilun Gemicioğlu⁵**, **Fuat Kalyoncu⁶**¹Department of Otorhinolaryngology, Medical Faculty, Eskisehir Osmangazi University, Eskisehir, Turkey²Department of Otorhinolaryngology, Medical Faculty, Kirikkale University, Kirikkale, Turkey³Department of Pulmonary Diseases, Medical Faculty, Celal Bayar University, Manisa, Turkey⁴ENT Department, Gungoren Private Hospital, Istanbul, Turkey⁵Department of Pulmonary Diseases, Medical Faculty, Cerrahpaşa University, İstanbul, Turkey⁶Department of Pulmonary Diseases, Medical Faculty, Hacettepe University, Ankara, Turkey**ABSTRACT**

OBJECTIVES. As a person ages, the mucosal epithelium degenerates over time and the elderly often have complications due to dehydration. Nasal airflow decreases and nasal airway resistance increases. In this review, we discuss the relationship of rhinitis, sinusitis and asthma in elderly patients.

MATERIAL AND METHODS. The literature survey was performed using the search engines PubMed, Google, Proquest Central, and EBSCO at the Kirikkale University Library.

RESULTS. With the aging process, allergic rhinitis decreases, whereas non-allergic rhinitis, such as drug-induced rhinitis can be observed more. Atrophic rhinitis is often seen in the elderly. Patients with geriatric rhinitis typically complain of constant sinus drainage problems, which may cause sinusitis. In addition, asthma prevalence has also been shown to be higher in elderly patients. Concurrent upper (rhinitis) and lower (asthma) respiratory tract symptoms are defined as Combined Allergic Rhinitis and Asthma Syndrome.

CONCLUSION. Rhinitis and asthma as well as sinusitis and asthma may be present in elderly patients. The incidence of allergic rhinitis has been shown to decrease with age. To achieve good results from treatment, the concurrent rhinitis and asthma as well as sinusitis and asthma should be kept in mind in elderly patients.

KEYWORDS: elderly, rhinitis, sinusitis, asthma, Combined Allergic Rhinitis and Asthma Syndrome, respiratory tract.

INTRODUCTION

Rhinitis is common in the elderly, and it is thought that a large amount of the general public has nasal symptoms¹⁻³. It is thought that the morphological changes in the connective tissue, as well as the vessel walls, predisposes elderly people to a more significant level of chronic rhinitis compared to the general population⁴.

Older people are known to have a lower whole

body water content, degenerative mucus-secreting glands, decreased mucociliary clearance. Furthermore, blood flow reduction to the nasal cavity promotes mucosal membrane drying and atrophy. On a structural basis, upper and lower nasal cartilage weakening and distortion of the nasal tip result in the resistance of the airway^{1,3}. All of these characteristics contribute to the nasal complications observed in the elderly.

In this review, we discuss the relationship of

Corresponding author: Dr. Nuray Bayar Muluk, Birlik Mahallesi, Zirvekent 2. Etap Sitesi, C-3 blok, No: 6-3/43, 06610, Çankaya / Ankara, Turkey

ORCID: <https://orcid.org/0000-0003-3602-9289>

e-mail: nuray.bayar@yahoo.com; nurayb@hotmail.com

Received for publication: 18 October 2021 / **Accepted:** 17 November 2021

rhinitis, sinusitis and asthma in elderly patients. The literature survey was performed using the search engines PubMed, Google, Proquest Central, and Ebsco at the Kırıkkale University Library.

CHANGES IN THE AGED NOSE

Nasal structure

As people grow older, various nasal anatomic changes can result in an increased incidence of rhinitis. This is thought to be associated with the loss of nasal tip support as one ages. A drooped tip is often caused by the erosion of collagen and elastin or facial musculature reduction⁵⁻⁷. These structural modification as one ages can lead to nasal airflow reduction and nasal obstruction, ultimately contributing to sinus complications.

Mucus

Previously published reports have shown that the mucosal epithelium deteriorates as the body ages and elderly patients have a tendency to dehydrate^{8,9}. Due to these factors, elderly patients frequently have an increase in mucus thickness. This characteristic combined with a reduction in mucociliary clearance has been thought to result in common nasal symptoms, such as cough and postnasal drip, all of which are associated with increased age⁵.

Nasal dryness

Clinically, it has been shown that the elderly are more prone to nasal dryness. Previous reports have shown that nasal cavity humidity and temperature were drastically lower in older patients compared to younger patients¹⁰. Interestingly, decreased humidity and temperature can be caused by age-related morphological changes to the nasal vasculature. For example, nasal vessels overtime can become patent and do not hydrate and sustain temperature as those of a younger nose⁹. These findings indicate that older patients experience a higher amount of nasal dryness and irritation due to these alterations.

Nasal airflow

Kalmovich and colleagues showed that endonasal volumes and cross-sectional regions progressively change with age¹¹. The reduction in estrogen content within the nasal mucosa contributes to hardening and increasing airway resistance. Due to the changes in estrogen levels, mucociliary modifications can also affect post-menopausal women¹². Recent reports have shown that certain estrogen receptors of the nasal mucosa associate with rhinitic symptoms¹³. However, the exact mechanism(s) has (have) not been defined.

Olfaction

It is known that the olfactory sense deteriorates

with age, and abundantly declines after the age of 70¹⁴. For this reason, the olfactory function over time may associate with rhinitis. One report showed that multiple study participants (71%) had a dysosmia and a positive allergy skin test association¹⁵. Olfactory complications in allergic rhinitis (AR) patients have generally shown to associate with nasal obstruction. However, recent data has shown that the etiology may be associated with olfactory cleft inflammation¹⁶.

RHINITIS IN ELDERLY

Rhinitis causes nasal mucosa inflammation and its symptoms include rhinorrhea, congestion, post-nasal drip, itchy nose and sneezing¹⁷. In older people, these symptoms are exacerbated by the excessive drainage, olfactory sensory loss and dryness^{5,18,19}.

The aging nose becomes dry for multiple reasons: 1 - dehydration; 2 - degenerative mucosal gland, and 3 - blood flow reduction in the nose. Isotonic saline lavage can be used to reduce nasal dryness and enhance mucosal clearance^{1,20}.

Rhinitis has been classified as follows: allergic and nonallergic rhinitis (AR and NAR)¹⁹. AR is an IgE-mediated inflammatory complication triggered by multiple allergens including dust, pollens or molds. Symptoms of AR can be considered seasonal or persistent. According to the Allergic Rhinitis and its Impact on Asthma (ARIA) guidelines, intermittent AR is defined as those symptoms that last less than four successive weeks. Persistent AR is when symptoms last greater than four successive weeks. Symptom severity is considered either 'mild' or 'moderate/severe' and both are based on interference with everyday tasks²¹.

NAR is a non-IgE-mediated complication as associated with congestion and clear rhinorrhea, but with a prominent reduction in other symptoms^{22,23}. The symptoms associated with NAR can be perennial or sporadic and may be triggered by food, odors, environmental changes, or even emotion^{21,24,25}. NAR can present itself in older patients and typical symptoms include atrophy, vasomotor impairment, gustatory and medication-related allergies^{4,26}.

Epidemiology

Interestingly, there is not a lot of research on the incidence of NAR when compared to AR in elderly patients. Approximately 19 million people suffer in the U.S. from NAR²⁷. The occurrence of NAR is higher in females and the frequency enhances by age²⁸⁻³⁰. More than sixty percent of patients over 50 are diagnosed with the NAR subtype^{19,25}.

Rhinitis on quality of life

Benninger and colleagues discovered, using the

Rhinosinusitis Disability Index (RSDI), that AR can cause sleep disturbance³¹. Poor sleep habits have been well documented in elderly³², and thus AR may vastly contribute to these complications. The poor sleeping habits can contribute to other crucial physiologic processes, including cognition, endocrine function, glucose metabolism and appetite control as well as lower self-reported health status, depression and increased mortality³³⁻³⁸. The effect of AR on these processes may hamper the ability to perform everyday tasks¹⁷.

Pathophysiology

The aging process can affect many physiological processes including T-cell responsiveness, lymphocyte response, cytokine production and receptor expression^{39,40}. Interestingly, this is a paradox due to the fact that the occurrence of AR declines as age increases. The reduction in immune response may be affected by the level of malignancy or infection in the elderly^{39,41}, and the cytokine bursts may clarify the questions concerning chronic rhinitis.

Other immune responses change with age as well^{39,42}. This may explain why the elderly are more susceptible to infection^{39,41,43}. These alterations may also result in the milder symptoms of AR as well as its decreased incidence in the older population.

Types of rhinitis in elderly

Allergic rhinitis

The peak occurrence of AR is 37 per 1,000 persons and it is reported between 10 and 15 years of age. The occurrence of AR in people over 65 years is 3 per 1,000 persons⁴⁴. In fact, the National Center for Health Statistics report (2005) indicated that people between 45-64 years, 65-75 years, and older than 75 were struck by AR at 10.7%, 7.8% and 5.4%⁴⁵. Along with the non-specific immune changes, anatomic and morphologic changes of the nose have been thought to contribute to AR in older people³⁹.

Non-allergic rhinitis (idiopathic, vasomotor)

Idiopathic rhinitis, or vasomotor rhinitis (VR), is defined as a persistent inflammation of the nasal membrane with no known allergy, infection, structural lesions or systemic disease^{1,46}.

VR is the most common form of NAR, and its occurrence increases with age²⁸. Environmental stimuli (e.g., ozone, cigarette smoke) cause nasal inflammation and the triggering of neurogenic reflexes. Sensory nerves within the nose respond to these factors, which initiates sneezing by reflex^{47,48}. VR in the elderly may be a deterioration in the regulation of neurological reactions on nasal physiology¹⁹.

Drug-induced rhinitis

Interestingly, greater than 400 brand name drugs have rhinitis as a secondary effect. Elderly

patients frequently administering these drugs for their ailments may result in drug-induced rhinitis. Because of this, older patients may be at risk due to thinning and dryness of the nasal mucosa¹.

A host of anti-hypertensive drugs may result in nasal obstruction and conjugated estrogens may also augment airway resistance¹.

Aspirin has been shown to trigger bronchospasm in those that have nasal polyps and asthma. However, it has been shown to be associated with severe rhinitis in asthmatics in the presence or absence of polyps¹.

Other drugs, like psychotropic drugs and Viagra, which are used by the elderly, can also contribute to rhinitis¹.

Non-allergic rhinitis with eosinophilia

Non-allergic rhinitis with eosinophilia (NARES) is caused by eosinophil infiltration in the nasal passage. Typical symptoms are perennial nasal congestion and rhinorrhea. This can usually be treated by a corticosteroid¹.

Gustatory rhinitis

Gustatory rhinitis is a condition consisting of profuse rhinorrhea that is elicited by eating highly seasoned foods. Additionally, cold air may exacerbate this condition. Anticholinergic drugs may be useful in gustatory rhinitis reduction. Ipratropium bromide is also effective and safe¹.

Gustatory rhinorrhea is caused by the ingestion of certain foods and results in watery rhinorrhea. These symptoms can be socially debilitating and result in decreased consumption of food and nutrition. Generally, alcohol and spicy or cold foods cause this condition. The capsaicin content in spicy foods elicits the response that triggers the parasympathetic nervous system⁴⁷.

Atrophic rhinitis

Atrophic rhinitis is often observed in the elderly and complications such as degeneration and/or crusting of the nasal mucous membrane occur⁴⁹. It may also be caused by organisms such as *Klebsiella ozaenae*, on rare occasions. In some cases, secondary forms of atrophic rhinitis may be caused by nasal surgery^{1,50}.

Primary atrophic rhinitis is often called geriatric rhinitis due to the age-related changes in nasal physiology⁵¹. Histopathological changes have been shown to associate with geriatric rhinitis⁵². Garcia and colleagues showed that these changes lead to excessive evaporation of the mucus layer and decreased mucosal surface area, which result in thickened mucus and altered nasal airflow⁵³. Recent reports have documented a role for apoptosis in rhinitis, and showed a decrease in caspase-3 activity, which is a key in the apoptotic cascade⁵⁴.

CHRONIC RHINOSINUSITIS IN ELDERLY

Chronic rhinosinusitis (CRS) is defined as an inflammatory condition of the nasal passages that continues for 12 weeks or longer⁵⁵.

There are four symptoms of CRS: mucopurulent drainage; nasal obstruction; facial pain, pressure and/or fullness; decreased sense of smell⁵⁶.

CRS consists of different disease entities. CRS without nasal polyps (CRSsNP) can be distinguished from CRS with nasal polyps (CRSwNP). CRSwNP patients have a higher total symptom score and generally suffer from nasal obstruction. CRSsNP have more headaches compared to CRSwNP patients⁵⁷⁻⁶⁰.

Elderly patients may have multiple systemic diseases associated with rhinosinusitis. Several factors such as age, duration of symptoms, length between follow-up, and diabetes mellitus were found to influence the complication rate. Interestingly, diabetes mellitus patients showed an approximate 3-fold higher complication rate than patients without. The elderly patients controlled for these factors had less complications⁶¹.

Patients with geriatric rhinitis generally had nagging sinus drainage, chronic throat clearing and nasal obstruction, which was exacerbated when lying down⁶².

Chronic rhinosinusitis patients typically have a longer history of thick foul-smelling drainage, nasal obstruction, headaches and facial pressure. Conversely, the geriatric rhinitis patients have a dry, irritated nose. Chronic rhinosinusitis diagnosis can be confirmed by a CT scan of the sinus cavity⁶².

ASTHMA IN ELDERLY

Asthma is prevalent in older people and is more frequent than previously believed. Interestingly, the morbidity and mortality rates for elderly asthmatics are higher than younger patients⁶³. Due to the aging of the world's population⁶⁴, the burden of asthma is expected to rise dramatically. However, the pathomechanisms of asthma in the older population have not been determined. Clinical observations indicate a role for several factors on adult or late-onset disease⁶³⁻⁶⁵. Several risk factors have been documented for elderly-associated asthma including smoking, obesity and SE-specific IgE sensitization⁶⁶⁻⁷⁰.

RHINITIS AND ASTHMA ASSOCIATION

AR and allergic asthma often exist together. This has led to the concept that these conditions are quite possibly the same complication. When patients have either AR or allergic asthma, they

are thoroughly examined. Often, observations of allergic inflammation and airway sensitivity throughout all of the airways is found⁷¹.

Non-allergic diseases of the nose can result in the advancement or worsening of asthma, mainly in children. Even in patients with NAR, an increased risk that asthma will develop is possible⁷¹.

Even though the upper airway and lungs share similarities, there are distinct differences. The upper airway and sinuses have large blood supply and, when inflamed (AR), can cause nasal blockage. On the other hand, the lungs respond to inflammation by spasming and become constricted and narrowed, causing chest tightness, wheezing and cough⁷¹.

Common colds and other viral infections are often associated with asthma worse outcomes. People with allergies are more susceptible to developing asthma after a viral respiratory tract infection⁷¹.

The upper airway performs as a multifaceted regulator of heat, humidity, resonance and filtration. Based on this, inhaled particles bigger than 5-6 microns are typically captured and retained in the upper airway⁷¹. This region of the lungs can elicit an immunologic response to allergens, which generally leads to inflammation of the upper and lower airways. Even though an allergic individual seems symptom free, continuous inflammation in the airway still exists⁷¹.

Upper or lower airway symptoms can be caused by specific allergens that an individual is exposed to. Larger allergens (~5 microns) such as grass or tree pollen, cause intermittent symptoms and are usually lodged in the nose and eye. Smaller antigens, such as animal dander or dust mites, are more likely to cause AR or asthma and are inhaled/captured in the lungs⁷¹.

Combined Allergic Rhinitis and Asthma Syndrome (CARAS) is defined as simultaneous upper (rhinitis) and lower (asthma) respiratory infection¹. Doctors suggest that patients with AR or sinusitis be tested for asthma due to the airway sensitivity⁷¹. Treatment of CARAS include:

- Topical corticosteroids can be used⁷¹.
- Oral administration such as systemic corticosteroids, anti-histamines and anti-leukotrienes⁷¹.
- Anti-IgE antibody therapy, which is a new class of anti-allergy⁷¹.
- Allergen immunotherapy is important in the CARAS treatment⁷¹.

SINUS DISEASE AND ASTHMA

Airway hyperresponsiveness in sinusitis is a result of the pharyngobronchial reflex activation⁷². This reflex may be exacerbated by infected drain-

age from sinuses that ooze into the throat⁷³. In patients with CRS, the pharyngeal mucosa becomes damaged and the pharyngeal nerve fiber density increases⁷⁴.

It has been well documented that sinus diseases elicit a plethora of inflammatory mediators such as eosinophils, lymphocytes and cytokines. These factors facilitate the recruitment of mast cells and basophils. Patients with AR generally have allergic inflammation. Furthermore, allergen induced activation of systemic responses can provoke inflammatory cell production and cause an augmentation of inflammatory cells that migrate to the airways⁷⁵⁻⁷⁸.

Interestingly, approximately 70% of patients indicated rhinosinusitis symptoms in a study comparing corticosteroid-dependent versus independent asthmatic patients⁷⁹. In another study, 24% of asthma and 36% of patients with small airway disease had CRS and were non-responders to medical treatment⁷⁹. Also, approximately seven percent have nasal polyposis⁸⁰, and the percentages are much more for nonatopic asthma patients (13%)⁸¹⁻⁸⁴.

CONCLUSIONS

Rhinitis and asthma, as well as sinusitis and asthma, are problematic in elderly patients. CARAS is the simultaneous upper and lower respiratory infection, which is related to nasal structural changes, nasal airflow decreases, and nasal obstruction often found in geriatric patients. The incidence of AR decreases with age, while drug induced rhinitis may be observed more in elderly patients and may be related to the drugs used for the treatment of geriatric diseases.

Conflict of interest: The author Cemal Cingi declares that he has no conflict of interest. The author Nuray Bayar Muluk declares that she has no conflict of interest. The author Arzu Yorgancıoğlu declares that she has no conflict of interest. The author Necdet Demir declares that he has no conflict of interest. The author Bilun Gemicioğlu declares that she has no conflict of interest. The author Fuat Kalyoncu declares that he has no conflict of interest.

Contribution of authors: Cemal Cingi - Planning, designing, literature survey. Nuray Bayar Muluk - Planning, designing, literature survey, writing. Arzu Yorgancıoğlu - Planning, designing, literature survey. Necdet Demir - Planning, designing, literature survey. Bilun Gemicioğlu - Planning, designing, literature survey. Fuat Kalyoncu - Planning, designing, literature survey.

Ethics committee approval: This is a review paper, there is no need for Ethics Committee approval.

Informed consent: There is no need to take informed consent because this is a review paper.

Founding: There is no funds for this article. There are no financial disclosures of the authors.

Acknowledgements: Preparation of this paper including design and planning was supported by Continuous Education and Scientific Research Association. There is no financial support. There is only scientific support.

REFERENCES

1. Slavin RG. Treating rhinitis in the older population: special considerations. *Allergy Asthma Clin Immunol.* 2009;5(1):9. DOI: 10.1186/1710-1492-5-9.
2. Hatter JB, Ouslander JG, Tinetti M, Studenski S, High KP, Asthana S. (eds.). *Hazzard's Geriatric Medicine and Gerontology.* 6 Edition. New York: McGraw Hill; 2009.
3. Sibbold B. Epidemiology of allergy rhinitis. In: Burr ML. (Editor). *Epidemiology of clinical allergy. Monographs in allergy.* Basel, Kar-gen; 1993, p. 61-79.
4. Slavin RG. Diagnosis and treatment of rhinitis and sinusitis in the elderly. *Immunol Allergy Clinics.* 1997;17(4):543-56. DOI: 10.1016/S0889-8561(05)70329-5.
5. Edelstein D. Aging of the normal nose in adults. *Laryngoscope.* 1996;106(9 Pt 2):1-25. DOI: 10.1097/00005537-199609001-00001.
6. Patterson C. The aging nose: characteristics and correction. *Otolaryngol Clin North Am.* 1980;13(2):275-88.
7. Reiss M, Reiss G. Rhinitis in old age. *Praxis (Bern 1994).* 2002;91(9):353-8. DOI: 10.1024/0369-8394.91.9.353.
8. Schrödter S, Biermann E, Halata Z. Histological evaluation of age-related changes in human respiratory mucosa of the middle turbinate. *Anat Embryo (Berl).* 2003;207:19-27. DOI: 10.1007/s00429-003-0326-5.
9. Bende M. Blood flow with ¹³³Xe in human nasal mucosa in relation to age, sex and body position. *Acta Otolaryngol.* 1983;96(1-2):175-9. DOI: 10.3109/00016488309132889.
10. Lindemann J, Sannwald D, Wiesmiller K. Age-related changes in intranasal air conditioning in the elderly. *Laryngoscope.* 2008;118(8):1472-5. DOI: 10.1097/MLG.0b013e3181758174.
11. Kalmovich LM, Elad D, Zaretsky U, Adunsky A, Chetrit A, Sadetzki S, et al. Endonasal geometry changes in elderly people: acoustic rhinometry measurements. *J Gerontol A Biol Sci Med Sci.* 2005;60(3):396-8. DOI: 10.1093/Gerona/60.3.396.
12. Nappi C, Sardo AS, Guerra G, Bifulco G, Testa D, Di Carlo C. Functional and morphologic evaluation of the nasal mucosa before and after hormone therapy in postmenopausal women with nasal symptoms. *Fertil Steril.* 2003;80(3):669-71. DOI: 10.1016/s0015-0282(03)00975-0.
13. Philpott CM, Wild DC, Wolstensholme CR, Murty GE. The presence of ovarian hormone receptors in the nasal mucosa and their relationship to nasal symptoms. *Rhinology.* 2008;46(3):221-5.
14. Seiberling KA, Conley DB. Aging and olfactory and taste function. *Otolaryngol Clin North Am.* 2004;37(6):1209-28. DOI: 10.1016/j.otc.2004.06.006.
15. Apter AJ, Mott AE, Frank ME, Clive JM. Allergic rhinitis and olfactory loss. *Ann Allergy Asthma Immunol.* 1995;75(4):311-6.

16. Pinto JM, Jeswani S, Reder L, Wang J, Perez R, DeTineo M, et al. Olfactory cleft inflammation is present in seasonal allergic rhinitis and improves with intranasal steroids abstract. *J Allergy Clin Immunol.* 2008;121(2 Suppl 1):S277. DOI: 10.1016/j.jaci.2007.12.1146.
17. Dykewicz MS, Fineman S, Skoner DP, Nicklas R, Lee R, Blessing-Moore J, et al. Diagnosis and management of rhinitis: complete guidelines of the Joint Task Force on Practice Parameters in Allergy, Asthma and Immunology. *Ann Allergy Asthma Immunol.* 1998;81(5 Pt 2):478-518. DOI: 10.1016/S1081-1206(10)63155-9.
18. Chadwick S. Allergic rhinitis in elderly. In: Calhoun K, Eibling D, editors. *Geriatric Otolaryngology.* New York: Taylor & Francis Group; 2006, p. 213-24.
19. Pinto JM, Jeswani S. Rhinitis in the geriatric population. *All Asth Clin Immun.* 2010;6(1):10. DOI: 10.1186/1710-1492-6-10.
20. Yilmaz ASS, Corey JP. Rhinitis in the elderly. *Curr Allergy Asthma Rep.* 2006;6:125-31. DOI: 10.1007/s11882-006-0050-3.
21. Bousquet J, Khaltaev N, Cruz AA, Denburg J, Fokkens WJ, Togias A, et al. Allergic Rhinitis and its Impact on Asthma (ARIA) 2008 update (in collaboration with the World Health Organization, GA(2)LEN and AllerGen). *Allergy.* 2008;63(Suppl 86):8-160. DOI: 10.1111/j.1398-9995.2007.01620.x.
22. Wheeler PW, Wheeler SF. Vasomotor rhinitis. *Am Fam Physician.* 2005;72(6):1057-62.
23. Bousquet J, Fokkens W, Burney P, Durham SR, Bachert C, Akdis CA, et al. Important research questions in allergy and related diseases: nonallergic rhinitis: a GA2LEN paper. *Allergy.* 2008;63(7):842-53. DOI: 10.1111/j.1398-9995.2008.01715.x.
24. Ayars G. Nonallergic rhinitis. *Immunol Allergy Clin North Am.* 2005;20(2):283-302. DOI: 10.1016/S0889-8561(05)70149-1.
25. Sanico A, Togias A. Noninfectious, nonallergic rhinitis (NINAR): considerations on possible mechanisms. *Am J Rhinol.* 1998;12(1):65-72. DOI: 10.2500/105065898782103043.
26. Yilmaz AAS, Corey JP. Rhinitis in the elderly. *Clin Allergy Immunol.* 2007;19:209-19.
27. Settupane RA, Charnock DR. Epidemiology of rhinitis: allergic and nonallergic. *Clin Allergy Immunol.* 2007;19:23-34.
28. Lal D, Corey JP. Vasomotor rhinitis update. *Curr Opin Otolaryngol Head Neck Surg.* 2004;12(3):243-7. DOI: 10.1097/01.moo.0000122310.13359.79.
29. Georgitis JW. Prevalence and differential diagnosis of chronic rhinitis. *Curr Allergy Asthma Rep.* 2001;1(3):202-6. DOI: 10.1007/s11882-001-0006-6.
30. Bachert C, van Cauwenberge P, Olbrecht J, van Schoor J. Prevalence, classification and perception of allergic and nonallergic rhinitis in Belgium. *Allergy.* 2006;61(6):693-8. DOI: 10.1111/j.1398-9995.2006.01054.x.
31. Benninger MS, Benninger RM. The impact of allergic rhinitis on sexual activity, sleep, and fatigue. *Allergy Asthma Proc.* 2009;30(4):358-65. DOI: 10.2500/aap.2009.30.3244.
32. Feinsilver SH. Sleep in the elderly. What is normal? *Clin Geriatr Med.* 2003;19(1):177-88. DOI: 10.1016/s0749-0690(02)00064-2.
33. Spiegel K, Leproult R, Van Cauter E. Impact of sleep debt on metabolic and endocrine function. *Lancet.* 1999;345(9188):1435-9. DOI: 10.106/S0140-6736(99)01376-8.
34. Trenell MI, Marshall NS, Rogers NL. Sleep and metabolic control: waking to a problem? *Clin Exp Pharmacol Physiol.* 2007;34(1-2):1-9. DOI: 10.1111/j.1440-1681.2007.04541.x.
35. Byles JE, Mishra GD, Harris MA, Nair K. The problems of sleep for older women: changes in health outcomes. *Age Ageing.* 2003;32(2):154-63. DOI: 10.1093/ageing/32.2.154.
36. Kim JM, Stewart R, Kim SW, Yang SJ, Shin I, Yoon J. Insomnia, depression, and physical disorders in late life: a 2-year longitudinal community study in Koreans. *Sleep.* 2009;32(9):1221-8. DOI: 10.1093/sleep/32.9.1221.
37. Zancocci M, Ponzetto M, Spada S, Rizzo R, Aimar T, Maero B, et al. Sleep disorders in the aged. *Minerva Med.* 1999;90(11-12):421-7.
38. Dew MA, Hoch CC, Buysse DJ, Monk TH, Begley AE, Houck PR, et al. Healthy older adults' sleep predicts all-cause mortality at 4 to 19 years of follow-up. *Psychosom Med.* 2003;65(1):63-73. DOI: 10.1097/01.psy.0000039756.23250.7c.
39. Bellanti JA, Azem M, MacDowell-Carneiro A, Tutuncuoglu SO, Wallerstedt DB. Possible mechanisms of late-life-onset allergic diseases and asthma in the senior citizen. *Allergy Asthma Proc.* 2000;21(5):267-70. DOI: 10.2500/108854100778248287.
40. Uciechowski P, Kahmann L, Plümäkers B, Malavolta M, Mocchegiani E, Dedoussis G, et al. TH1 and TH2 cell polarization increases with aging and is modulated by zinc supplementation. *Exp Gerontol.* 2008;43(5):493-8. DOI: 10.1016/j.exger.2007.11.006.
41. Hwang KA, Kim HR, Kang I. Aging and human CD4(+) regulatory T cells. *Mech Ageing Dev.* 2009;130(8):509-17. DOI: 10.1016/j.mad.2009.06.003.
42. Gruver AL, Hudson LL, Sempowski GD. Immunosenescence of ageing. *J Pathol.* 2007;211(2):144-56. DOI: 10.1002/path.2104.
43. Ostan R, Bucci L, Capri M, Salvioli S, Scurti M, Pini E, et al. Immunosenescence and immunogenetics of human longevity. *Neuroimmunomodulation.* 2008;15(4-6):224-40. DOI: 10.1159/000156466.
44. Pederson PA, Weeke ER. Allergic rhinitis in Danish general practice. Prevalence and consultation rates. *Allergy.* 1981;36(6):375-9. DOI: 10.1111/j.1398-9995.1981.tb01841.x.
45. Mahmoudi M. Prevalance of allergic diseases in children, adults, and the elderly In: Shanahan J, Naglieri C, editors. *Asthma and Allergy: Practical Diagnosis and Management.* 1. McGraw-Hill Professional; 2008, p. 18-20.
46. Lund VJ, Aaronson A, Bousquet J, Dahl R, Davies RJ, Durham SR, et al. International consensus report on the diagnosis and management of rhinitis. *Allergy.* 1994;49(Suppl 19):2-34.
47. Salib RJ, Harries PG, Nair SB, Howarth PH. Mechanisms and mediators of nasal symptoms in non-allergic rhinitis. *Clin Exp Allergy.* 2008;38(3):393-404. DOI: 10.1111/j.1365-2222.2007.02926.x.
48. Shi G, Li X. The roles of three peptides in the pathogenesis of vasomotor rhinitis. *Lin Chuang Er Bi Yan Hou Ke Za Zhi.* 2006;20(17):777-9, 783.
49. Liston SL, Siegel LG. Nasal and sinus disorders in the elderly: which ones are life-threatening? *Geriatrics.* 1981;36(2):91-102.
50. Moore GF, Freeman TJ, Ogren FP, Yonkers AJ. Extended follow-up of total inferior turbinate resection for relief of chronic nasal obstruction. *Laryngoscope.* 1985;95(9 Pt 1):1095-9.
51. Jordan JA, Mabry RL. Geriatric rhinitis: what it is, and how to treat it. *Geriatrics.* 1998;53(76):81-4.
52. Mehrotra R, Singhal J, Kawatra M, Gupta SC, Singh M. Pre and post-treatment histopathological changes in atrophic rhinitis. *Indian J Pathol Microbiol.* 2005;48(3):310-3.
53. Garcia GJM, Bailie N, Martins DA, Kimbell JS. Atrophic rhinitis: a CFD study of air conditioning in the nasal cavity. *J Appl Physiol.* (1985). 2007;103(3):1082-92. DOI: 10.1152/jappphysiol.01118.2006.
54. Hirt R, Paulsen F, Neumann K, Knipping S. Immunocytochemical detection of caspase 3 in various diseases of human nasal mucosa. *HNO.* 2009;57(5):466-72. DOI: 10.1007/s00106-009-1905-4.
55. Scadding GK, Durham SR, Mirakian R, Jones NS, Drake-Lee AB, Ryan D, et al. BSACI guidelines for the management of rhinosinusitis and nasal polyposis. *Clin Exp Allergy.* 2008;38(2):260-75. DOI: 10.1111/j.1365-2222.2007.02889.x.
56. Hamilos DL, Holbrook EH. Chronic rhinosinusitis: Management.

- [Internet]. UpToDate. 2016. Available from: <https://www.uptodate.com/contents/chronic-rhinosinusitis-management#H1>.
57. Fokkens W, Lund V, Mullol J. European Position Paper on Rhinosinusitis and Nasal Polyps group. European position paper on rhinosinusitis and nasal polyps 2007. *Rhinol Suppl.* 2007;20:1-136.
 58. Bachert C, Claeys SEM, Tomassen P, van Zele T, Zhang N. Rhinosinusitis and asthma: a link for asthma severity. *Curr Allergy Asthma Rep.* 2010;10(3):194-201. DOI: 10.1007/s11882-010-0096-0.
 59. Huvenne W, van Bruaene N, Zhang N, van Zele T, Patou J, gevaert P, et al. Chronic rhinosinusitis with and without nasal polyps: what is the difference? *Curr Allergy Asthma Rep.* 2009;9(3):213-20. DOI: 10.1007/s11882-009-0031-4.
 60. Bachert C, Van Bruaene N, Toskala E, Zhang N, Olze H, Scadding G, et al. Important research questions in allergy and related diseases: 3-chronic rhinosinusitis and nasal polyposis – a GALEN study. *Allergy.* 2009;64(4):520-33. DOI: 10.1111/j.1398-9995.2009.01964.x.
 61. Ban JH, Kwon HJ, Lee KC. Outcomes of endoscopic sinus surgery in an elderly population: comparison with those in an adult population. *Clin Otolaryngol.* 2010;35(4):300-6. DOI: 10.1111/j.1749-4486.2010.02155.x.
 62. American Anademy of Otolaryngology-Head and Neck Surgery. Sinusitis: Special considerations for aging patients. [Internet]. Available from: <https://coastalear noseandthroat.com/wp-content/uploads/2020/12/Sinusitis-Special-Considerations.pdf>.
 63. Yáñez A, Cho SH, Soriano JB, Rosenwasser IJ, Rodrigo GJ, Rabe KF, et al. Asthma in the elderly: what we know and what we have yet to know. *World Allergy Organ J.* 2014;7(1):8. DOI: 10.1186/1939-4551-7-8.
 64. Christensen K, Doblhammer G, Rau R, Vaupel JW. Ageing populations: the challenges ahead. *Lancet.* 2009;374(9696):1196-1208. DOI: 10.1016/S0140-6736(09)61460-4.
 65. Hanania NA, King MJ, Braman SS, Saltoun C, Wise RA, Enright P, et al. Asthma in the elderly: Current understanding and future research needs—a report of a National Institute on Aging (NIA) workshop. *J Allergy Clin Immunol.* 2011;128(3 Suppl):S4-24. DOI: 10.1016/j.jaci.2011.06.048.
 66. Song WJ, Kim SH, Lim S, Park YJ, Kim MH, Lee SM, et al. Association between obesity and asthma in the elderly population: potential roles of abdominal subcutaneous adiposity and sarcopenia. *Ann Allergy Asthma Immunol.* 2012;109(4):243-8. DOI: 10.1016/j.anai.2012.07.017.
 67. Kim YK, Kim SH, Tak YJ, Jee YK, Lee BJ, Kim SH, et al. High prevalence of current asthma and active smoking effect among the elderly. *Clin Exp Allergy.* 2002;32(12):1706-12. DOI: 10.1046/j.1365-2222.2002.01524.x.
 68. Song WJ, Sintobin I, Sohn KH, Kang MG, Park HK, Jo EJ, et al. Staphylococcal enterotoxin IgE sensitization in late-onset severe eosinophilic asthma in the elderly. *Clin Exp Allergy.* 2016;46(3):411-21. DOI: 10.1111/cea.12652.
 69. Bachert C, van Steen K, Zhang N, Holtappels G, Cattaert T, Maus B, et al. Specific IgE against *Staphylococcus aureus* enterotoxins: an independent risk factor for asthma. *J Allergy Clin Immunol.* 2012;130(2):376-81.e8. DOI: 10.1016/j.jaci.2012.05.012.
 70. Tomassen P, Jarvis D, Newson R, Van Ree R, Forsberg B, Howarth P, et al. *Staphylococcus aureus* enterotoxin-specific IgE is associated with asthma in the general population: a GA(2)LEN study. *Allergy.* 2013;68(10):1289-97. DOI: 10.1111/all.12230.
 71. Ferreira LKDP, Ferreira LAMP, Monteiro TM, Bezerra GC, Bernardo LR, Puivezam MR. Combined allergic rhinitis and sthma syndrome (CARAS). *Int Immunopharmacol.* 2019;74:105718. DOI: 10.1016/j.intimp.2019.105718.
 72. Smart BA, Slavlin RG. Rhinosinusitis and pediatric asthma. *Immunol Allergy Clin North Am.* 2005;25(1):67-82. DOI: 10.1016/j.ia.2004.09.002.
 73. Bachert C, Patou J, Van Cauwenberge P. The role of sinus disease in asthma. *Curr Opin Allergy Clin Immunol.* 2006;6(1):29-36. DOI: 10.1097/01.all.0000200504.54425.0e.
 74. Rolla G, Colagrande P, Scappaticci E, Bottomicca F, Magnano M, Brussino L, et al. Damage of the pharyngeal mucosa and hyperresponsiveness of airway in sinusitis. *J Allergy Clin Immunol.* 1997;100(1):52-7. DOI: 10.1016/s0091-6749(97)70194-5.
 75. Braunstahl GJ, Kleinjan A, Overbeek SE, Prins JB, Hoogsteden HC, et al. Segmental bronchial provocation induces nasal inflammation in allergic rhinitis patients. *Am J Respir Crit Care Med.* 2000;161(6):2051-7. DOI: 10.1164/ajrccm.161.6.9906121.
 76. Stirling RG, van Rensen EL, Barnes PJ, Chung KF. Interleukin-5 induces CD34(+) eosinophil progenitor mobilization and eosinophil CCR3 expression in asthma. *Am J Respir Crit Care Med.* 2001;164(8 Pt 1):1403-9. DOI: 10.1164/ajrccm.164.8.2010002.
 77. Dorman SC, Efthimiadis A, Babirad I, Watson RM, Denburg JA, Hargreave FE, et al. Sputum CD34+IL-5Ralpha+ cells increase after allergen: evidence for in situ eosinophilopoiesis. *Am J Respir Crit Care Med.* 2004;169(5):573-7. DOI: 10.1164/rccm.200307-1004OC.
 78. Braunstahl GJ, Overbeek SJ, Kleinjan A, Prins JB, Hoogsteden HC, Fokkens WJ. Nasal allergen provocation induces adhesion molecule expression and tissue eosinophilia in upper and lower airways. *J Allergy Clin Immunol.* 2001;107(3):469-76. DOI: 10.1067/mai.2001.113046.
 79. Bresciani M, Paradis L, Des Roches A, Vernhet H, Vachier I, Godard P, et al. Rhinosinusitis in severe asthma. *J Allergy Clin Immunol.* 2001;107(9):73-80. DOI: 10.1067/mai.2001.111593.
 80. Ragab A, Clement P, Vincken W. Objective assessment of lower airway involvement in chronic rhinosinusitis. *Am J Rhinol.* 2004;18(1):15-21.
 81. Settupane GA, Chafee FH. Nasal polyps in asthma and rhinitis: a review of 6,037 patients. *J Allergy Clin Immunol.* 1977;59(1):17-21. DOI: 10.1016/0091-6749(77)90171-3.
 82. Settupane GA. Epidemiology of nasal polyps. In: Settupane GA, Lund VJ, Bernstein JM, Tos M (eds.). *Nasal polyps: epidemiology, pathogenesis and treatment.* Providence, Rhode Island: Oceanside Publications; 1997, p. 17-24.
 83. Lamblin C, Gosset P, Salez F, Vandezande LM, Perez T, Darras J, et al. Eosinophilic airway inflammation in nasal polyposis. *J Allergy Clin Immunol.* 1999;104(1):85-92. DOI: 10.1016/s0091-6749(99)70118-1.
 84. Larsen K. The clinical relationship of nasal polyps to asthma. *Allergy Asthma Proc.* 1996;17(5):243-49. DOI: 10.2500/108854196778662255.

