

## EDITORIAL

# Aerosol therapy in ENT disorders

Raluca Enache , MD, PhD

ENT Sarafoleanu Medical Clinic, Bucharest, Romania



Aerosols are microparticles of a liquid or solid nature dispersed in a gaseous medium. Aerosol therapy refers to the administration of certain medicinal substances in the form of microparticles to the upper respiratory tract. Initially conceived as a method of systemic treatment (due to the large absorption surface at the pulmonary level), this therapeutic method is now primarily used for its topical effect on the respiratory mucosa. The benefits of aerosol therapy have been recognized since ancient times<sup>1-3</sup>. The first reference to the use of aerosols as a treatment method dates back to 1554 BC, recorded on an Egyptian papyrus found in the Theban necropolis<sup>4</sup>. Their use in the treatment of asthma was described as early as 600 BC by two Indian physicians, Charaka and Sushruta, who used various herbs with anticholinergic properties<sup>2</sup>. In ancient Greece, Hippocrates (460-377 BC) and, later, Galen of Pergamon (2<sup>nd</sup> century AD) described the use of therapeutic aerosols for various conditions, including nasal, pulmonary, and laryngeal pathologies<sup>1,5</sup>.

Primarily used in the treatment of bronchopulmonary diseases, aerosol-based therapy is also particularly useful in numerous ENT conditions<sup>6</sup>. In 2005, the NUAGES survey, which monitored the use of nebulizers in the medical field in France, revealed that this therapy was most frequently prescribed by pulmonologists, with ENT specialists ranking second at that time, accounting for 89% of prescriptions<sup>7,8</sup>.

A significant group of sinonasal conditions can benefit from aerosol therapy. Aerosols have been successfully used in the treatment of rhinosinusitis or rhinitis, including allergic rhinitis<sup>6,9-12</sup>. The main advantage of the therapy is the high concentration of the substances used at the level of the rhinosinus mucosa, leading to

a rapid therapeutic effect and minimal systemic response. Larger aerosol particles most frequently deposit in the upper airways, with their concentration increasing with particle size (90% for 10 $\mu$ m particles, and 10% for 2 $\mu$ m particles)<sup>3,7,13,14</sup>. Particles smaller than 5 $\mu$ m tend to deposit more in the mucosa of the oral cavity, larynx, and lungs<sup>3,13</sup>. The accumulation of particles in the sinonasal mucosa is also influenced by the air flow velocity; the higher the velocity, and thus the speed of particle movement, the greater the level of deposition in the upper airways.

There are different types of aerosol generators that can be used: sprays or nebulizers. Sprays release particles with sizes ranging from 10 to 150 $\mu$ m, at a high velocity, resulting in a greater accumulation in the anterior region of the nasal cavities<sup>7</sup>. Nebulizers release particles at a lower velocity, ranging from 1 to 10 $\mu$ m, thus favoring their deposition also in the posterior region of the nasal cavities<sup>7</sup>, thereby increasing the treatment's efficacy. Among all types of nebulizers, there are studies indicating that pulsating aerosol devices may be more effective in delivering a larger amount of medication to the paranasal sinuses<sup>6,9,15</sup>.

Aerosol therapy represents an extremely useful therapeutic method in combination with oral or intravenous medication. Depending on the sinonasal pathology, different medications can be used in aerosol therapy: saline solutions, corticosteroids, mucolytics, antibiotics, and even essential oils. Essential oils, such as eucalyptus or menthol, improve mucociliary clearance and have antibacterial, antifungal, and antiviral effects<sup>16,17</sup>. The use of corticosteroids in aerosols, although controversial and insufficiently studied, seems to have slightly greater efficacy in reducing rhinosi-

**Corresponding author:** Raluca Enache, MD, PhD, ENT Sarafoleanu Medical Clinic, 1 Lt. Av. Iuliu Tetrat Street, District 1, Bucharest, Romania

**ORCID:** <https://orcid.org/0000-0002-2841-6265>

**e-mail:** r.enache@rinologie.ro

**Received for publication:** September 10, 2024 / **Accepted:** September 18, 2024

nusitis inflammatory symptoms compared to topical steroid nasal sprays<sup>9</sup>. Administration should not exceed 10 days of treatment. Antibiotics delivered locally through aerosols can accumulate in higher concentrations in the rhinosinus mucosa, providing evident benefits<sup>9</sup>. There are studies showing a positive effect of this type of treatment in the case of rhinosinusitis refractory to medical and surgical treatment<sup>7,9,10</sup>.

Aerosol-based treatment in conjunction with antibiotic and anti-inflammatory therapy can also be found in the therapeutic protocol for acute or chronic laryngitis and vocal nodules.

This therapeutic method is also useful in the treatment of Eustachian tube dysfunction or chronic otitis media with effusion, in which case the aerosols are made from a mixture of vasoconstrictor and/or antibiotic substances. The sessions performed have a beneficial effect on the mucosa of the Eustachian tube and the middle ear.

Due to the wide range of medicinal substances used in aerosol therapy (anti-inflammatory, mucolytic, vasoconstrictor, antibiotic), the therapeutic effects can vary.

Regarding aerosol devices, there are three types of devices used: pressurized metered-dose inhalers (MDIs) primarily used in patients with pulmonary conditions, dry powder inhalers (DPIs) and nebulizers.

In ENT pathology, nebulizers are the most frequently used devices. They do not require patient coordination and are used to deliver a larger amount of medication in a short time. They utilize high quantities of medicinal substances, leading to increased concentrations at the mucosal and pulmonary levels. The high operating pressure ensures a high nebulization rate for greater therapeutic efficacy. Smaller-capacity nebulizers can be used for both adults and children, even during sleep.

Particularly effective as a method of administration, under the right indication, aerosol-based therapy represents a therapeutic alternative in inflammatory or infectious pathologies within the field of ENT.

**Funding:** None.

**Conflict of interests:** None.

**Financial discloser:** None.

## REFERENCES

- Stein SW, Thiel CG. The history of therapeutic aerosols: a chronological review. *J Aerosol Med Pulmon Drug Del.* 2017;30(1):20-41. DOI: 10.1089/jamp.2016.1297.
- Rospond B, Krakowska A, Muszynska B, Opoka W. The history, current state and perspective of aerosol therapy. *Acta Pharm.* 2022;72:225-43. DOI: 10.2478/acph-2022-0017.
- Amirav I. Aerosol therapy. *Italian Journal of Pediatrics.* 2004;30(3):147-56.
- Breasted JH. *The Edwin Smith surgical papyrus.* University of Chicago Press, Chicago, IL; 1930.
- Shehata M. History of inhalation therapy. *Internet J Health.* 2008;9:1-9.
- Durand M, Le Guellec S, Pourchez J, Dubois F, Aubert G, Chantel G, et al. Sonic aerosol therapy to target maxillary sinuses. *Eur Ann Otorhinolaryngol Head Neck Dis.* 2012;129(5):244-50. DOI: 10.1016/j.ano.2011.09.002.
- Pruliere-Escabasse V, Michel J, Percodani J, Serrano E, Gilain L, Crampette L., et al. Consensus document for prescription of nebulization in rhinology. *Eur Ann Otorhinolaryngol Head Neck Dis.* 2014;131(6):371-4. DOI: 10.1016/j.ano.2014.07.004.
- De Monte M, Dubus JC, Chaumuzeau JP, Dautzenberg B, Dessanges JF, Becquemain MH, et al. A survey of nebulization practices in France conducted in 2004 by the GAT. *Rev Mal Respir* 2008;25(1):43-9. DOI: 10.1016/s0761-8425(08)70465-0.
- Veleplic M, Manestar D, Perkovic I, Skalamera D, Braut T. Inhalation aerosol therapy in the treatment of chronic rhinosinusitis: a prospective randomized study. *J Clin Pharmacol.* 2019;59(12):1648-55. DOI: 10.1002/jcph.1471.
- Shikani AH, Khoueir N, Jabra-Rizk MA, Shikani HJ, Basaraba RJ, Leid JG. Topical therapy for refractory rhinosinusitis caused by methicillin-resistant *Staphylococcus aureus*: First report in a prospective study. *Auris Nasus Larynx.* 2018;45(5):994-9. DOI: 10.1016/j.anl.2018.01.009.
- Ohki M, Hyo Y, Yoshiyama Y, Takano H, Takahata J, Suzuki M, et al. Consensus guidance of nebulizer therapy for acute rhinosinusitis. *Auris Nasus Larynx.* 2020;47(1):18-24. DOI: 10.1016/j.anl.2019.08.007.
- Fokkens WJ, Lund VJ, Hopkins C, Hellings PW, Kern R, Reitsma S, et al. European Position Paper on Rhinosinusitis and nasal Polyps 2020. *Rhinology.* 2020;58(Suppl S29):1-464. DOI: 10.4193/Rhin20.600.
- Fry FA. Charge distribution on polystyrene aerosols and deposition in human nose. *Aerosol Sci* 1970:195.
- Badre R, Guillermin R. Diffusion et rétention des aérosols thérapeutiques dans les voies respiratoires supérieures. *Poumon.* 1979;35:341-7.
- Moller W, Schuschnig U, Celik G, Munzing W, Bartenstein P, Haussinger K, et al. Topical drug delivery in chronic rhinosinusitis patients before and after sinus surgery using pulsating aerosols. *PLoS One.* 2013;8(9):e74991. DOI: 10.1371/journal.pone.0074991.
- Elaissi A, Rouis Z, Salem NA, Mabrouk S, Salem YB, Salah KBH, et al. Chemical composition of 8 eucalyptus species' essential oils and the evaluation of their antibacterial, antifungal and antiviral activities. *BMC Complement Altern Med.* 2012;12:81. DOI: 10.1186/1472-6882-12-81.
- Lai Y, Dilidaer D, Chen B, Xu G, Shi J, Lee RJ, et al. In vitro studies of a distillate of rectified essential oils on sinonasal components of mucociliary clearance. *Am J Rhinol Allergy.* 2014;28(3):244-8. DOI: 10.2500/ajra.2014.28.4036.



This is an open access article published under the terms and conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>). CC BY-NC-ND 4.0 license requires that reusers give credit to the creator by citing or quoting the original work. It allows reusers to copy, share, read, download, print, redistribute the material in any medium or format, or to link to the full texts of the articles, for non-commercial purposes only. If others remix, adapt, or build upon the material, they may not distribute the modified material.